
CHAPTER 13

IPM FOR SCORPIONS IN SCHOOLS

INTRODUCTION

Scorpions live in a wide variety of habitats from tropical to temperate climates and from deserts to rain forests. In the United States they are most common in the southern states from the Atlantic to the Pacific. All scorpions are beneficial because they are predators of insects.

The sting of most scorpions is less painful than a bee sting. There is only one scorpion of medical importance in the United States: the sculptured or bark scorpion, *Centruroides exilicauda* (=sculpturatus). The danger from its sting has been exaggerated, and its venom is probably not life-threatening. This species occurs in Texas, western New Mexico, Arizona, northern Mexico, and sometimes along the west bank of the Colorado River in California.

IDENTIFICATION AND BIOLOGY


Scorpions range from 3/8 to 8 1/2 inches in length, but all scorpions are similar in general appearance.

Scorpions do not lay eggs; they are viviparous, which means they give birth to live young. As embryos, the young receive nourishment through a kind of “placental” connection to the mother’s body. When the young are born, they climb onto the mother’s back where they remain from two days to two weeks until they molt (shed their skin) for the first time. After the first molt, the young disperse to lead independent lives. Some scorpions mature in as little as six months while others take almost seven years.

All scorpions are predators, feeding on a variety of insects and spiders. Large scorpions also feed on small animals including snakes, lizards, and rodents. Some scorpions sit and wait for their meal to come to them while others actively hunt their prey. Scorpions have a very low metabolism and some can exist for 6 to 12 months without food. Most are active at night. They are shy creatures, aggressive only toward their prey. Scorpions will not sting humans unless handled, stepped on, or otherwise disturbed.

It is rare for scorpions to enter a building since there is little food and temperatures are too cool for their comfort. There are some exceptions to this rule. Buildings

in new developments (less than three years old) can experience an influx of scorpions because the construction work has destroyed the animals’ habitat. In older neighborhoods, the heavy bark on old trees provide good habitat for scorpions, and they may enter through the more numerous cracks and holes in buildings in search of water, mates, and prey. Also, buildings near washes and arroyos that are normally dry may become refuges for scorpions during summer rains.



Scorpions do not enter buildings in winter because cold weather makes them sluggish or immobile. They are not active until nighttime low temperatures exceed 70°F. Buildings heated to 65½ or 70°F provide enough warmth to allow scorpions to move about. Scorpions found inside buildings in cold weather are probably summer visitors that never left. Although scorpions prefer to live outdoors, they can remain in buildings without food for long periods of time.

STINGS

A scorpion sting produces considerable pain around the site of the sting, but little swelling. For four to six hours, sensations of numbness and tingling develop in the region of the sting, then symptoms start to go away. In the vast majority of cases, the symptoms will subside within a few days without any treatment.

If the sting is from a bark scorpion, symptoms can sometimes travel along nerves, and tingling from a sting on a finger may be felt up to the elbow, or even the shoulder. Severe symptoms can include roving eyes, blurry vision, excessive salivation, tingling around the mouth and nose, and the feeling of having a lump in the throat. Respiratory distress may occur. Tapping the sting can produce extreme pain. Symptoms in children also include extreme restlessness, excessive muscle activity, rubbing at the face, and sometimes vomiting. Most vulnerable to the sting of the bark scorpion are children under five years and elderly persons who have

Avoiding Scorpion Stings

Schools in areas where encounters with scorpions are likely should teach children and adults how to recognize scorpions and to understand their habits. Focus on scorpion biology, behavior, likely places to find them, and how to avoid disturbing them.

At home, children and parents should be taught to take the following precautions to reduce the likelihood of being stung:

- Wear shoes when walking outside at night. If scorpions are suspected indoors, wear shoes inside at night as well.
- Wear leather gloves when moving rocks, boards, and other debris.
- Shake out shoes or slippers before they are worn, and check beds before they are used.
- Shake out damp towels, washcloths, and dishrags before use.
- When camping, shake out sleeping bags, clothes, and anything else that has been in contact with the ground before use.
- Protect infants and children from scorpions at night by placing the legs of their cribs or beds into clean wide-mouthed glass jars and moving the crib or bed away from the wall. Scorpions cannot climb clean glass.

an underlying heart condition or respiratory illness. The greatest danger to a child is the possibility of choking on saliva and vomit.

Antivenin for the bark scorpion is produced at Arizona State University in Tempe and is available in Arizona but not in other states. The therapeutic use of antivenin is still experimental. People have been treated without antivenin for many years, and in areas where antivenin is unavailable, people are monitored closely by medical professionals until the symptoms subside.

DETECTION AND MONITORING

To determine where scorpions are entering, inspect both the inside and outside of the building at night (when scorpions are active) using a battery-operated camp

light fitted with a UV (black) fluorescent bulb. Scorpions glow brightly in black light and can be spotted several yards away.

Always wear leather gloves when hunting scorpions. Places to check inside the building include under towels, washcloths, and sponges in bathrooms and kitchen; under all tables and desks, since the bark scorpion may climb and take refuge on a table leg or under the lip of a table; and inside storage areas. Outside, check piles of rocks and wood, under loose boards, and in piles of debris. After the following treatment strategies have been implemented, monitoring with the black light should continue to verify population reduction.

MANAGEMENT OPTIONS

Physical Controls

In most cases, physical controls will be adequate to manage a scorpion problem.

Removal of Scorpions

Any scorpions found during monitoring can be picked up using gloves or a pair of kitchen tongs, and transferred to a clean, wide-mouthed glass jar. Scorpions cannot climb clean glass. You can also invert a jar over a live scorpion and then slide a thin piece of cardboard under the mouth of the jar to trap the scorpion inside. Once a scorpion is captured, drop it into a jar of alcohol or soapy water (water without soap will not work) to kill it.

Habitat Modification

If you discover areas near school buildings that harbor a number of scorpions, you can try to alter the habitat to discourage them. Wood piles, rocks, loose boards, and other debris should be removed from the immediate vicinity of the building.

If there are slopes on the school grounds that are faced with rip rap (large rocks placed on a slope or stream bank to help stabilize it), or other similar areas highly attractive to scorpions, place a barrier of aluminum flashing between the rip rap and the school to prevent scorpion access. The flashing must be bent in an “L” shape away from the building. The other edge of the flashing should be buried a short distance from the rocks, deep enough in the soil so that the L shape will not fall over and lean on the rip rap. Make the height of the barrier before the bend greater than the height of the rip rap to prevent scorpions from standing on the rocks and jumping over the barrier.

Carry a caulking gun during nighttime inspections inside and outside the building to seal any cracks and crevices found. If scorpions are entering through weep holes in windows or sliding doors, cover the holes with fine-mesh aluminum screening, available from hardware or lumber stores. The ends of pipes that are designed as gray water drains should be fitted with loose filter bags, or makeshift end-pieces made from window screen. The screened end will prevent scorpion access to the drainpipe, sink, and other parts of the building.

It is important to continue nighttime patrols and caulking until all entryways have been located and sealed, and all the scorpions in the building have been captured and killed. Once the access routes are sealed, and all indoor scorpions have been removed, only doorways provide access, unless the scorpions ride in on logs and other materials. Glazed tiles can be placed around the perimeter of the buildings, and under or around doors and windows as part of the decor and as practical scorpion barriers. Scorpions have difficulty crossing smooth tiles unless the grout line is wide. Wood storage should be elevated above the ground since scorpions like contact with moist soil. Before bringing materials such as logs inside, bang them on a stone to dislodge any scorpions.

Traps

A simple trap made of damp gunny sacks laid down near the building in the evenings may be useful for monitoring and trapping. Scorpions may seek out the moist environment under the sacks where they can be collected in the morning. This trap is most effective when used before summer rains.

Chemical Controls

In general, preventing scorpion problems is better than trying to kill these creatures with pesticide. Spraying the perimeters of buildings is not only unnecessary but also ineffective. Scorpions can tolerate a great deal of pesticide in their environment. The pesticide will only be harmful to humans, especially children, and to other wildlife. Using physical controls along with education to reduce the fear of scorpions will help prevent unnecessary treatments.

First Aid for Scorpion Stings

Most scorpion stings are similar to a bee or wasp sting, and like bee or wasp stings, the majority of scorpion stings can be treated at school or the victim's home.

First aid for a scorpion sting includes the following:

- Calm the victim.
- Do not use a tourniquet.
- Wash the area with soap and water.
- Apply a cool compress (an ice cube wrapped in a wet washcloth), but do not apply ice directly to the skin or submerge the affected limb in ice water.
- To reduce pain, over-the-counter pain relievers such as aspirin or acetaminophen can help.
- Elevate or immobilize the affected limb if that feels more comfortable.
- Do not administer sedatives such as alcohol.

BIBLIOGRAPHY

- Bio-Integral Resource Center (BIRC). 1996. 1997 directory of least-toxic pest control products. *IPM Practitioner* 18(11/12):1-39.
- Cloudsley-Thompson, J.L. 1968. *Spiders, Scorpions, Centipedes and Mites*. Pergamon Press, New York. 278 pp.
- Keegan, H.L. 1980. *Scorpions of Medical Importance*. Jackson University Press, Jackson, MS. 140 pp.
- Mallis, A. 1982. *Handbook of Pest Control*. Franzak and Foster, Cleveland, OH. 1101 pp.
- Olkowski, W., S. Daar, and H. Olkowski. 1991. *Common-Sense Pest Control: Least-toxic solutions for your home, garden, pets and community*. Taunton Press, Newtown, CT. 715 pp.
- Polis, G.A., ed. 1990. *The Biology of Scorpions*. Stanford University Press, Stanford, CA. 587 pp.
- Smith, R.L. 1982. *Venomous animals of Arizona*. Cooperative Extension Service, College of Agriculture, University of Arizona, Tucson, AZ, Bulletin 8245. 134 pp.
- Smith, R.L. 1992. Personal communication. Associate Professor, Entomology Dept., University of Arizona at Tucson.

